


The background of the slide is a photograph of two common frog tadpoles (Knoblauchkröte) in a clear plastic container. The tadpoles are submerged in water, surrounded by green grass and some dry plant matter. They have brown and black mottled patterns on their bodies. The text is overlaid on this image.

Erste Resultate zur Wiederansiedlung der Knoblauchkröte im Lüsekamp und Elmpter Schwalmbruch

Stefanie Stablum

Biologische Station Krickenbecker Seen

Die Knoblauchkröte (*Pelobates fuscus*)
Knoflookpad


















KK-Start



Flächenvorstellung

- 3 geeignete Gewässer
- Sandiger Boden
- Fischfrei
- Tackenbenden
- Lüsekamp Trichter
- Lüsekamp Nord







Tackenbenden

2020

250 Larven
25 Jungkröten



Lüsekamp Nord

250 Larven
75 Jungkröten



Lüsekamp Trichter

30 Larven
25 Jungkröten



Tackenbenden



2021

500 Larven
40 Jungkröten



Lüsekamp Nord



500 Larven
39 Jungkröten




Lüsekamp Trichter




100 Larven
31 Jungkröten

Erfassung Tackenbenden



Erfassung Lüsekamp Nord



Erfassung Lüsekamp Trichter

1

■ Männchen ■ Weibchen

Projekt Niederlande

- Eigenes Zuchtprogramm
- 2011 Fang letzter Exemplare Rondven
- Aussetzung 2011-2015, 2017
- Seit 2014 3 Flächen (Plateau)
- Seit 2017 4 Flächen (Amphibienpoel)
- Individualerkennung



Ergebnisse: Plateau

Jahr	Individuen
2015	21
2022	1


Ergebnisse: Rondven

Jahr	Individuen
2022	10

Ergebnisse: Amphibienpoel


Jahr	Individuen
2017	75
2018	72
2022	28

Gesamt 2022



Ausblick

?





Fragen?